

# Folic Acid Supplementation to Prevent Neural Tube Defects

## Updated Evidence Report and Systematic Review for the US Preventive Services Task Force

Meera Viswanathan, PhD; Rachel Peragallo Urrutia, MD, MS; Kesha N. Hudson, PhD; Jennifer Cook Middleton, PhD; Leila C. Kahwati, MD, MPH

**IMPORTANCE** Neural tube defects are among the most common birth defects in the US.

**OBJECTIVE** To review new evidence on the benefits and harms of folic acid supplementation for the prevention of neural tube defects to inform the US Preventive Services Task Force.

**EVIDENCE REVIEW** Sources included PubMed, Cochrane Library, Embase, and trial registries from July 1, 2015, through July 2, 2021; references; and experts, with surveillance through February 10, 2023. Two investigators independently reviewed English-language randomized studies and nonrandomized cohort studies in very highly developed countries that focused on the use of folic acid supplementation for the prevention of neural tube defect–affected pregnancies; methodological quality was dually and independently assessed.

**FINDINGS** Twelve observational studies (reported in 13 publications) were eligible for this limited update ( $N = 1244\,072$ ). Of these, 3 studies ( $n = 990\,372$ ) reported on the effect of folic acid supplementation on neural tube defects. For harms, 9 studies were eligible: 1 randomized clinical trial ( $n = 431$ ) reported on variations in twin delivery, 7 observational studies ( $n = 761\,125$ ) reported on the incidence of autism spectrum disorder, and 1 observational study ( $n = 429\,004$ ) reported on maternal cancer. Two cohort studies and 1 case-control study newly identified in this update reported on the association between folic acid supplementation and neural tube defects ( $n = 990\,372$ ). One cohort study reported a statistically significant reduced risk of neural tube defects associated with folic acid supplementation taken before pregnancy (adjusted relative risk [aRR], 0.54 [95% CI, 0.31-0.91]), during pregnancy (aRR, 0.62 [95% CI, 0.39-0.97]), and before and during pregnancy (aRR, 0.49 [95% CI, 0.29-0.83]), but this association occurred for only the later of 2 periods studied (2006-2013 and not 1999-2005). No other statistically significant benefits were reported overall. No study reported statistically significant harms (multiple gestation, autism, and maternal cancer) associated with pregnancy-related folic acid exposure.

**CONCLUSIONS AND RELEVANCE** New evidence from observational studies provided additional evidence of the benefit of folic acid supplementation for preventing neural tube defects and no evidence of harms related to multiple gestation, autism, or maternal cancer. The new evidence was consistent with previously reviewed evidence on benefits and harms.

JAMA. 2023;330(5):460-466. doi:10.1001/jama.2023.9864

← Editorial pages 417 and 419

+ Multimedia

← Related article page 454 and JAMA Patient Page page 480

+ CME at [jamacmlookup.com](#)

**Author Affiliations:** RTI International—University of North Carolina at Chapel Hill Evidence-based Practice Center (Viswanathan, Hudson, Middleton, Kahwati); RTI International, Research Triangle Park, North Carolina (Viswanathan, Hudson, Kahwati); Department of Obstetrics and Gynecology, University of North Carolina at Chapel Hill (Urrutia); Cecil G. Sheps Center for Health Services Research, University of North Carolina at Chapel Hill (Middleton).


**Corresponding Author:** Meera Viswanathan, PhD, RTI International, 3040 E Cornwallis Rd, PO Box 12194, Research Triangle Park, NC 27709 ([viswanathan@rti.org](mailto:viswanathan@rti.org)).

**N**eural tube defects are major congenital malformations often caused by low folate concentrations in the body at the time of conception. These defects frequently result in significant disability or death for affected fetuses and children. Strategies that enhance folic acid uptake before pregnancy offer the best chance of prevention.

In 2017, the US Preventive Services Task Force (USPSTF) concluded that folic acid supplementation in the periconceptional period has substantial benefits in reducing the risk of neural tube defects in the developing fetus<sup>1</sup> and reaffirmed its 2009 recommendation that all persons who are planning or capable of preg-

nancy take a daily supplement containing 0.4 to 0.8 mg (400-800 µg) of folic acid (A recommendation). The 2017 USPSTF recommendation was based on previously reviewed evidence from a randomized clinical trial and observational studies reporting reduced neural tube defects with supplementation and no consistent evidence of harms such as multiple gestation, maternal adverse effects, or child respiratory illness.

This limited evidence update aimed to identify studies published since the previous (2017) evidence review<sup>2</sup> conducted for the USPSTF to inform a reaffirmation of the current recommendation.

**Figure. Analytic Framework: Folic Acid Supplementation to Prevent Neural Tube Defects**

Evidence reviews for the US Preventive Services Task Force (USPSTF) use an analytic framework to visually display the key questions that the review will address to allow the USPSTF to evaluate the effectiveness and safety of a preventive service. The questions are depicted by linkages that relate to interventions and outcomes. Further details are available from the USPSTF procedure manual.<sup>3</sup> NTD indicates neural tube defect.

## Methods

An analytic framework and 2 key questions guided the limited evidence update (Figure). A literature search of PubMed/MEDLINE, the Cochrane Library, Embase, and trial registries was conducted from July 1, 2015, through July 2, 2021. Additional sources included reference lists of retrieved articles, outside experts, and public commenters, with ongoing surveillance of the literature through February 10, 2023. Two investigators independently evaluated the eligibility of all abstracts and articles and rated study quality using predefined criteria.<sup>4</sup> Detailed methods and results are available in the full evidence report.<sup>4</sup>

English-language randomized and nonrandomized studies that focused on the use of folic acid supplementation (by itself or in multivitamins) for the prevention of neural tube defect-affected pregnancies in persons capable of getting pregnant were eligible. Studies conducted in very highly developed countries and that investigated potential harms of folic acid supplementation, such as maternal cancer and autism spectrum disorder, were also eligible. Ineligible studies included poor-quality studies and those focusing solely on persons taking antiseizure medications or with a history of neural tube defects in previous pregnancies.

## Results

Twelve observational studies (reported in 13 publications<sup>5-17</sup>) (Table) were eligible for this limited update ( $N = 1244\ 072$  [from nonoverlapping cohorts]). Of these, 3 studies ( $n = 990\ 372$ ) assessed the effect of folic acid supplementation on neural tube defects.<sup>5-8</sup> No studies examined differences by race or ethnicity.

For harms, 9 studies were eligible; 1 randomized clinical trial ( $n = 431$ ) assessed variations in twin delivery,<sup>9</sup> 7 observational studies ( $n = 761125$ ) examined the incidence of autism spectrum disorders,<sup>10-16</sup> and 1 observational study ( $n = 429\ 004$ ) reported on maternal cancer.<sup>17</sup> The Table also reports details on studies from the 2017 evidence review.<sup>18-41</sup>

### Benefits of Folic Acid Supplementation

Regarding the benefits of folic acid supplementation, 2 cohort studies and 1 case-control study in this update examined the association between folic acid supplementation and neural tube defects ( $n = 990\ 372$ ).<sup>5-8</sup> Food fortification and supplementation practices varied by setting. Of these studies, 1 cohort study set in Norway (no mandatory fortification) reported on neural tube defects among live births and stillborn infants from 1999 to 2013 overall and also stratified results into 2 separate periods: 1999 to 2005 and 2006 to 2013.<sup>6</sup> The authors performed this stratified analysis because they found that the overall adjusted relative risk (aRR) was affected by year of birth. Several external events of importance were cited to explain differences by period: the introduction of folic acid recommendations in 1999, inclusion of 0.2 mg of folic acid in multivitamin supplements from 2004 onward (before 2004, most multivitamins did not include folic acid), and increased adherence to folic acid recommendations in the second half of the period analyzed (2006-2013).<sup>6</sup> The authors reported no statistically significant benefits in the first of the 2 periods (1999-2005), regardless of timing of supplementation (before pregnancy, during pregnancy, or before and during pregnancy). In contrast, in the second period (2006-2013), the authors reported a statistically significant reduced risk of neural tube defects associated with folic acid supplementation taken before pregnancy (aRR, 0.54 [95% CI, 0.31-0.91]), during

Table. Summary of Evidence From Limited Update Review: Folic Acid Supplementation to Prevent Neural Tube Defects

Key question	Evidence summary in 2017 <sup>2</sup>	Summary of new evidence	Limitations of new evidence	Consistency of new evidence with prior evidence findings
KQ1: Benefits of folic acid supplementation				
1a: Effects of folic acid supplements on risk of NTDs	12 Studies (1 RCT, 2 cohort, 8 case-control, 1 previous review); n > 41 802 Generally consistent evidence within the prefortification (indicating benefit) and postfortification (no statistically significant differences) eras; inconsistent over time	3 Studies (2 cohort [3 publications <sup>5-7</sup> ], 1 case-control <sup>8</sup> ); N = 990 372 Norwegian cohort (no mandatory fortification) study reported no statistically significant associations in overall analysis (1999-2013) or the first period (1999-2005) with low adherence to folic acid supplementation recommendations	Heterogeneous populations with different levels of food fortification and diet patterns; methodological limitations in foundational evidence also apply	New studies have some evidence of benefit for reducing NTDs and do not change conclusions from foundational evidence
1. RCT (prefortification): Peto OR for NTD, 0.131 (95% CI, 0.026-0.648); P = .01 <sup>9,42-47</sup>		Statistically significant associations from 2006-2013 with higher adherence to folic acid recommendations: Before pregnancy only (aRR, 0.54 [95% CI, 0.31-0.91]) <sup>6</sup> During pregnancy only (aRR, 0.62 [95% CI, 0.39-0.97]) <sup>7</sup> Before and during pregnancy (aRR, 0.49 [95% CI, 0.29-0.83]) <sup>7</sup>		
2. Cohort studies (prefortification): aOR for NTD, 0.11 (95% CI, 0.01-0.91) <sup>18</sup>		No consistently and statistically significant associations in Japanese cohort of general population (no mandatory fortification) (aOR, 0.62 [95% CI, 0.23-1.71] for preconceptual use when compared with use after pregnancy recognition or no use) <sup>5</sup> or US and Canadian case-control study (postfortification study) of participants with prepregnancy diabetes or pregestational obesity for exposures measured as less than daily, daily, <0.4 mg, 0.4 mg to <1.0 mg <sup>8</sup>		
3. Case-control studies (prefortification): aOR for NTD, 0.7 (95% CI, 0.5-0.8) <sup>21</sup>				
RR for NTD, 0.6 (95% CI, 0.4-0.8) <sup>22</sup>				
OR for NTD, 0.65 (95% CI, 0.45-0.94) <sup>23</sup>				
OR for NTD, 1.00 (95% CI, 0.73-1.40); P = .97 <sup>24</sup>				
1. Case-control study (spanning prefortification and postfortification eras): aOR for NTD, 1.12 (95% CI, 0.22-5.78) <sup>25</sup>				
3. Case-control studies (postfortification): OR for NTD, 1.11 (95% CI, 0.74-1.65) for consistent users <sup>26</sup>				
aOR for NTD (anencephaly + spina bifida), 0.93 (95% CI, 0.82-1.06) <sup>27</sup>				
aOR (anencephaly), 1.2 (95% CI, 0.8-1.9) <sup>28</sup>				
aOR (spina bifida), 1.4 (95% CI, 1.0-1.8) <sup>28</sup>				
No new trials can be conducted on this topic				
New studies must rely on observational data with inherent risks of case ascertainment bias (in prospective cohort studies) or exposure recall bias (in retrospective studies)				
1b: Differences in effect of folic acid supplements on NTDs by race or ethnicity				
1. Differences in effect of folic acid supplements on NTDs by race or ethnicity	No effect in first study <sup>28</sup>	No new evidence	NA	NA
2. Differences in effect of folic acid supplements on NTDs by race or ethnicity	Higher risk in second study (aOR for Hispanic women with consistent use compared with nonuse, 2.20 [95% CI, 0.98-4.92]) <sup>26</sup>			
3. Differences in effect of folic acid supplements on NTDs by race or ethnicity	Less protective effect in third study (OR, 0.96 [95% CI, 0.44-2.10] for Hispanic women vs 0.62 [95% CI, 0.35-1.10] for non-Hispanic White women vs 0.54 [95% CI, 0.09-3.20] for Black women) <sup>23</sup>			
	Small numbers in each comparison, differences in direction of estimate of effects possibly due to chance			

(continued)

Table. Summary of Evidence From Limited Update Review: Folic Acid Supplementation to Prevent Neural Tube Defects (continued)

Key question	Evidence summary in 2017 <sup>2</sup>	Summary of new evidence	Limitations of new evidence	Consistency of new evidence with prior evidence findings
1c: Differences in effect of folic acid supplements on NTDs by dosage, duration, and timing	<p>Dosage: 4 studies (1 cohort, 3 case-control); n = 26 791 No indication of dose response in 3 of 4 studies;<sup>19,20,22-24</sup>, 1 study showed lower odds for daily use vs less than daily use (OR, 0.57 [95% CI, 0.35-0.93])<sup>22</sup></p> <p>Duration: 0 studies Timing: 5 studies (1 cohort, 4 case-control); n = 26 808 Calculated OR from cohort study for use weeks 1-6 vs weeks 7 and later, 0.29 (95% CI, 0.14-0.60)<sup>19,20</sup></p> <p>Older studies consistently showed no effect of timing<sup>23,25</sup>; 1 new study (postfortification) showed a protective effect of use before pregnancy vs initiation in the first month of pregnancy on anencephaly but not spina bifida<sup>28</sup>; the other new study did not find a protective effect for spina bifida for consistent periconceptional use vs initiation in the first month of pregnancy<sup>26</sup></p> <p>Small numbers in each comparison, effects possibly due to chance, studies used different measures of dose and timing</p>	<p>Dosage: 1 case-control study of women with pre pregnancy obesity; n = 1429 Statistically significantly reduced association between NTD risk and exposure of 0.4 mg to &lt;1.0 mg of folic acid supplementation daily (aOR, 0.54 [95% CI, 0.29-0.95]) but not for exposures of &lt;0.4 mg (aOR, 1.29 [95% CI, 0.40-3.37]) or &lt;0.4 mg or ≥1.0 mg (aOR, 0.84 [95% CI, 0.38-1.68]); differences did not persist in sensitivity analysis<sup>5</sup></p> <p>Duration: 0 studies Timing: 1 cohort study (2 publications<sup>6,7</sup>); n = 896 674 Consistent benefits regardless of timing in 1 of 3 periods examined (2006-2013) (aRR before pregnancy only, 0.54 [95% CI, 0.31-0.91]<sup>6</sup>; during pregnancy only, 0.62 [95% CI, 0.39-0.97]<sup>7</sup>; and before and during pregnancy, 0.49 [95% CI, 0.29-0.83])<sup>7</sup>; consistently no statistically significant differences for the other periods [1999-2013, 1999-2005]<sup>7</sup></p>	<p>Small numbers in each comparison, effects possibly due to chance, studies used different measures of dose and timing</p>	<p>New studies do not change conclusions regarding dosage or timing</p>
KQ2: Harms of folic acid supplementation				
2a: Harms associated with folic acid supplements; multiple gestation (twinning)	<p>2 Studies (1 trial, 1 cohort); n = 7387 Trial found no statistically significant differences in twin pregnancy rate (RR, 1.4 [95% CI, 0.87-2.26])<sup>29</sup></p> <p>Cohort study found higher risk of twin birth for folate use (OR, 1.59 [95% CI, 1.41-1.78]) that was attenuated once potential misclassification was accounted for (OR, 1.04 [95% CI, 0.91-1.18])<sup>30</sup></p> <p>Low event rate, wide CIs</p>	<p>No new evidence</p>	<p>NA</p>	<p>NA</p>
2a: Harms associated with folic acid supplements: childhood asthma, allergy, wheezing	<p>3 Systematic reviews, 8 observational studies; n &gt; 14 438 No effect for a large majority of comparisons and outcomes<sup>31-41</sup></p> <p>Variable measures of outcomes and exposure, all observation studies with risks of bias from case ascertainment and recall</p>	<p>No new evidence</p>	<p>NA</p>	<p>NA</p>
2a: Harms associated with folic acid supplements; other adverse events in women	<p>1 RCT; n = 4862 Increased risk for weight gain, diarrhea, constipation; reduced risk for irregular defecation; no difference for increased appetite, lack of appetite, exanthema, heartburn, and vertigo<sup>42</sup></p> <p>Low event rate, wide CIs</p>	<p>No new evidence</p>	<p>NA</p>	<p>NA</p>
2a: Harms associated with folic acid supplements; autism	<p>No eligible evidence</p>			
				<p>7 Studies (6 fair-quality cohort<sup>10-15</sup> and 1 fair-quality case-control<sup>16</sup>); n = 76 1125 Studies set in 4 countries (Israel, Sweden, Denmark, Norway); varied measures of exposure, comparators, and outcomes; generally no statistically significant associations; 3 publications on a populations in Israel<sup>15</sup> and Norway,<sup>12,14</sup> respectively, reported some benefits</p>

(continued)

Table. Summary of Evidence From Limited Update Review: Folic Acid Supplementation to Prevent Neural Tube Defects (continued)

Key question	Evidence summary in 2017 <sup>2</sup>	Summary of new evidence	Limitations of new evidence	Consistency of new evidence with prior evidence findings
2a: Harms associated with folic acid supplements: maternal cancer	No eligible evidence	1 Cohort study, <sup>17</sup> n = 429 004 HR for 1 pregnancy with exposure to folic acid supplementation vs no exposure in pregnancy, 1.08 (95% CI, 1.00–1.18) <sup>17</sup> HR for ≥2 pregnancies with exposure to folic acid supplementation vs no exposure in pregnancy, 1.06 (95% CI, 0.91–1.22) <sup>17</sup>	Potential for unmeasured confounding and recall bias in the classification of the intervention	NA
2b: Differences in harms associated with folic acid supplements by dosage, timing, and duration: twinning	No eligible evidence	1 Trial <sup>9</sup> ; n = 431 RR, 0.45 (95% CI, 0.11–1.77) for twin deliveries with exposure to 4-mg folic acid supplementation vs exposure to 0.4-mg folic acid supplementation; both groups exposed before conception and through 12 weeks of gestation	Applicability uncertain to unplanned pregnancies	NA
2b: Differences in harms associated with folic acid supplements by dosage, timing, and duration: childhood asthma, allergy, wheezing	Dosage: 1 systematic review, 1 observational study; n = 484 No consistent increase in the risk of childhood asthma, wheezing, or allergies by timing <sup>32,41</sup> Duration: 0 studies Timing: 2 systematic reviews, 3 observational studies; N varies by outcome No consistent increase in the risk of childhood asthma, wheezing, or allergies by timing <sup>31,33,34,40,41</sup> Variable measures of outcomes and exposure, all observational studies with risks of bias from case ascertainment and recall	No new evidence	NA	NA
2b: Differences in harms associated with folic acid supplements by dosage, timing, and duration: autism	No eligible evidence	Dosage: 3 studies (2 cohort, <sup>11,12</sup> 1 case-control) <sup>16</sup> ; n = 194 281 Overlap in CIs with exposure to folic acid supplementation in different doses vs no or very low exposure to folic acid supplementation in pregnancy, all not statistically significant Duration: 0 studies Timing: 2 cohort studies, <sup>11,12</sup> n = 120 235 Overlap in CIs with exposure to folic acid supplementation in different time intervals vs no exposure to folic acid supplementation in pregnancy, all but 1 estimate not statistically significant; initiation in weeks 5 to 8 associated with benefit (1.4/16 1.84 vs 32/14 7.21; aOR, 0.44 [95% CI, 0.23–0.83]) <sup>12</sup>	Potential for unmeasured confounding and recall bias in the classification of the intervention	NA

Abbreviations: aOR, adjusted odds ratio; aRR, adjusted relative risk; HR, hazard ratio; IQ, key question; NTD, neural tube defect; OR, odds ratio; RCT, randomized clinical trial; RR, relative risk.

pregnancy (aRR, 0.62 [95% CI, 0.39-0.97]), and before and during pregnancy (aRR, 0.49 [95% CI, 0.29-0.83]).<sup>6</sup>

The second cohort study, set in Japan (no mandatory food fortification), reported no statistically significant differences associated with adequate (preconception) folic acid supplementation (adjusted odds ratio [aOR], 0.62 [95% CI, 0.23-1.71]) when compared with inadequate use (use after pregnancy recognition or no use).<sup>5</sup> The third study, a case-control study set in the US and Canada in the period following food fortification, reported on participants with pregestational diabetes and prepregnancy obesity.<sup>8</sup> The study reported that cases occurred more often among persons with unplanned pregnancies.<sup>8</sup> Authors reported a statistically significant reduction in neural tube defects in women with prepregnancy obesity taking 0.4 mg to 1 mg of folic acid, when compared with women taking no supplementation and adjusting for maternal age (aOR, 0.54 [95% CI, 0.29-0.95]).<sup>8</sup> Results adjusting for planned pregnancy rather than maternal age were similar but not statistically significant (aOR, 0.57 [95% CI, 0.30-1.02]).<sup>8</sup> Across all 3 studies, no other statistically significant benefits were reported overall or by dose (1 study<sup>8</sup>) or timing (1 study<sup>6,7</sup>).

#### ARTICLE INFORMATION

**Accepted for Publication:** June 19, 2023.

**Author Contributions:** Dr Viswanathan had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

**Concept and design:** Viswanathan, Urrutia, Kahwati. **Acquisition, analysis, or interpretation of data:** Viswanathan, Urrutia, Hudson, Middleton.

**Drafting of the manuscript:** Viswanathan, Urrutia, Hudson, Middleton.

**Statistical analysis:** Viswanathan.

**Obtained funding:** Viswanathan, Kahwati.

**Administrative, technical, or material support:** Viswanathan, Hudson, Middleton, Kahwati.

**Supervision:** Viswanathan.

**Conflict of Interest Disclosures:** None reported.

**Funding/Support:** This research was funded under contract 75Q80120D00007, Task Order 01, from the Agency for Healthcare Research and Quality (AHRQ), US Department of Health and Human Services, under a contract to support the USPSTF.

**Role of the Funder/Sponsor:** Investigators worked with USPSTF members and AHRQ staff to develop the scope, analytic framework, and key questions for this review. AHRQ had no role in study selection, quality assessment, or synthesis. AHRQ staff provided project oversight, reviewed the report to ensure the analysis met methodological standards, and distributed the draft for peer review.

Otherwise, AHRQ had no role in the conduct of the study; collection, management, analysis, and interpretation of the data; and preparation, review, or approval of the manuscript findings. The opinions expressed in this document are those of the authors and do not represent the official position of AHRQ or the US Department of Health and Human Services.

**Additional Contributions:** We thank the following individuals for their contributions to this project: Justin Mills, MD, MPH (AHRQ medical officer); Tina Fan, MD, MPH (previous associate scientific director, AHRQ); and Tracy Wolff, MD, MPH (scientific director, AHRQ USPSTF program);

current and former members of the USPSTF; peer and federal partner reviewers; RTI International-University of North Carolina Evidence-based Practice Center staff: Christiane Voisin, MLS (research librarian); Roberta Wines, MPH, and Carol Woodell, BSPH (current and former Evidence-based Practice Center program managers); Nila Sathe, MA, MLIS (quality assurance); Sharon Barrell, MA (editor); and Teyonna Downing and Alex Cone (publications specialists). USPSTF members, peer reviewers, and federal partner reviewers did not receive financial compensation for their contributions.

**Additional Information:** A draft version of the full evidence report underwent external peer review from 3 content experts (Nancy Rose, MD, University of Utah; Jorge Chavarro, MD, ScD, Harvard University; Kimberly Gregory, MD, MPH, Cedars-Sinai Medical Center) and 3 individuals from 2 federal partners (Centers for Disease Control and Prevention, National Institutes of Health). Comments from reviewers were presented to the USPSTF during its deliberation of the evidence and were considered in preparing the final evidence review.

**Editorial Disclaimer:** This evidence report is presented as a document in support of the accompanying USPSTF recommendation statement. It did not undergo additional peer review after submission to *JAMA*.

#### REFERENCES

1. US Preventive Services Task Force. Folic acid supplementation for the prevention of neural tube defects: US Preventive Services Task Force recommendation statement. *JAMA*. 2017;317(2):183-189. doi:10.1001/jama.2016.19438
2. Viswanathan M, Treiman KA, Kish-Doto J, Middleton JC, Coker-Schwimmer EJL, Nicholson WK. *Folic Acid Supplementation: An Evidence Review for the US Preventive Services Task Force. Evidence Synthesis No. 145*. Agency for Healthcare Research and Quality; 2017. AHRQ publication 14-05214-EF-1.
3. US Preventive Services Task Force Procedure Manual. US Preventive Services Task Force. Published May 2021. Accessed July 11, 2023. <https://uspreventiveservicestaskforce.org/uspstf/about-uspstf/methods-and-processes/procedure-manual>
4. Viswanathan M, Urrutia RP, Hudson KN, Middleton JC, Kahwati LC. *Folic Acid Supplementation to Prevent Neural Tube Defects: A Limited Systematic Review Update for the US Preventive Services Task Force. Evidence Synthesis No. 230*. Agency for Healthcare Research and Quality; 2023. AHRQ publication 22-05302-EF-1.
5. Nishigori H, Obara T, Nishigori T, et al; Japan Environment & Children's Study Group. Preconception folic acid supplementation use and the occurrence of neural tube defects in Japan: a nationwide birth cohort study of the Japan Environment and Children's Study. *Congenit Anom (Kyoto)*. 2019;59(4):110-117. doi:10.1111/cga.12293
6. Gildestad T, Øyen N, Klungsøy K, Nilsen RM, Daltveit AK, Vollset SE. Maternal use of folic acid supplements and infant risk of neural tube defects in Norway 1999-2013. *Scand J Public Health*. 2016;44(6):619-626. doi:10.1177/1403494816649494
7. Gildestad T, Bjørge T, Haaland ØA, Klungsøy K, Vollset SE, Øyen N. Maternal use of folic acid and multivitamin supplements and infant risk of birth defects in Norway, 1999-2013. *Br J Nutr*. 2020;124(3):316-329. doi:10.1017/S0007114520001178
8. Petersen JM, Parker SE, Benedum CM, Mitchell AA, Tinker SC, Werler MM. Periconceptional folic acid and risk for neural tube defects among higher risk pregnancies. *Birth Defects Res*. 2019;111(19):1501-1512. doi:10.1002/bdr2.1579
9. Bortolus R, Filippini F, Cipriani S, et al. Efficacy of 4.0 mg versus 0.4 mg folic acid supplementation on the reproductive outcomes: a randomized controlled trial. *Nutrients*. 2021;13(12):4422. doi:10.3390/nu13124422
10. Strøm M, Granström C, Lyall K, Ascherio A, Olsen SF. Research letter: folic acid supplementation and intake of folate in pregnancy

#### Harms of Folic Acid Supplementation

No study of harms (multiple gestation, autism, and maternal cancer) reported significant associations with pregnancy-related folic acid exposure.<sup>9-17</sup>

#### Discussion

This evidence review identified 3 new observational studies reporting on the association between folic acid supplementation before or during pregnancy and neural tube defects in offspring. Mandatory food fortification and supplementation practices varied by geography and period of investigation and contributed to heterogeneity across studies. Nevertheless, these new studies provided additional evidence of the benefit of folic acid supplementation for preventing neural tube defects. Nine new observational studies found no evidence of harms related to multiple gestation, autism, or maternal cancer. This new evidence is consistent with previously reviewed evidence on the benefits and harms of folic acid supplementation to prevent neural tube defects.

- in relation to offspring risk of autism spectrum disorder. *Psychol Med.* 2018;48(6):1048-1054. doi:10.1017/S0033291717002410
11. Virk J, Liew Z, Olsen J, Nohr EA, Catov JM, Ritz B. Periconceptional and prenatal supplementary folic acid and multivitamin intake and autism spectrum disorders. *Autism.* 2016;20(6):710-718. doi:10.1177/1362361315604076
  12. Surén P, Roth C, Bresnahan M, et al. Association between maternal use of folic acid supplements and risk of autism spectrum disorders in children. *JAMA.* 2013;309(6):570-577. doi:10.1001/jama.2012.155925
  13. DeVilbiss EA, Magnusson C, Gardner RM, et al. Antenatal nutritional supplementation and autism spectrum disorders in the Stockholm youth cohort: population based cohort study. *BMJ.* 2017;359:j4273. doi:10.1136/bmj.j4273
  14. Nilsen RM, Surén P, Gunnes N, et al. Analysis of self-selection bias in a population-based cohort study of autism spectrum disorders. *Paediatr Perinat Epidemiol.* 2013;27(6):553-563. doi:10.1111/ppe.12077
  15. Levine SZ, Kodesh A, Viktorin A, et al. Association of maternal use of folic acid and multivitamin supplements in the periods before and during pregnancy with the risk of autism spectrum disorder in offspring. *JAMA Psychiatry.* 2018;75(2):176-184. doi:10.1001/jamapsychiatry.2017.4050
  16. Sharman Moser S, Davidovitch M, Rotem RS, Chodick G, Shalev V, Koren G. High dose folic acid during pregnancy and the risk of autism: the birth order bias: a nested case-control study. *Reprod Toxicol.* 2019;89:173-177. doi:10.1016/j.reprotox.2019.07.083
  17. Mortensen JH, Øyen N, Formina T, et al. Supplemental folic acid in pregnancy and maternal cancer risk. *Cancer Epidemiol.* 2015;39(6):805-811. doi:10.1016/j.canep.2015.10.009
  18. Czeizel AE, Dobó M, Varga P. Hungarian cohort-controlled trial of periconceptional multivitamin supplementation shows a reduction in certain congenital abnormalities. *Birth Defects Res A Clin Mol Teratol.* 2004;70(11):853-861. doi:10.1002/bdra.20086
  19. Milunsky A, Jick H, Jick SS, et al. Multivitamin/folic acid supplementation in early pregnancy reduces the prevalence of neural tube defects. *JAMA.* 1989;262(20):2847-2852. doi:10.1001/jama.262.20.2847
  20. Moore LL, Bradlee ML, Singer MR, Rothman KJ, Milunsky A. Folate intake and the risk of neural tube defects: an estimation of dose-response. *Epidemiology.* 2003;14(2):200-205. doi:10.1097/01.ede.0000040253.12446.b2
  21. Hernández-Díaz S, Werler MM, Walker AM, Mitchell AA. Neural tube defects in relation to use of folic acid antagonists during pregnancy. *Am J Epidemiol.* 2001;153(10):961-968. doi:10.1093/aje/153.10.961
  22. Werler MM, Shapiro S, Mitchell AA. Periconceptional folic acid exposure and risk of recurrent neural tube defects. *JAMA.* 1993;269(10):1257-1261.
  23. Shaw GM, Schaffer D, Velie EM, Morland K, Harris JA. Periconceptional vitamin use, dietary folate, and the occurrence of neural tube defects. *Epidemiology.* 1995;6(3):219-226. doi:10.1097/00001648-199505000-00005
  24. Mills JL, Rhoads GG, Simpson JL, et al; National Institute of Child Health and Human Development Neural Tube Defects Study Group. The absence of a relation between the periconceptional use of vitamins and neural-tube defects. *N Engl J Med.* 1989;321(7):430-435. doi:10.1056/nejm198908173210704
  25. Suarez L, Hendricks KA, Cooper SP, Sweeney AM, Hardy RJ, Larsen RD. Neural tube defects among Mexican Americans living on the US-Mexico border: effects of folic acid and dietary folate. *Am J Epidemiol.* 2000;152(11):1017-1023. doi:10.1093/aje/152.11.1017
  26. Ahrens K, Yazdy MM, Mitchell AA, Werler MM. Folic acid intake and spina bifida in the era of dietary folic acid fortification. *Epidemiology.* 2011;22(5):731-737. doi:10.1097/EDE.0b013e3182227887
  27. Agopian AJ, Tinker SC, Lupo PJ, Canfield MA, Mitchell LE; National Birth Defects Prevention Study. Proportion of neural tube defects attributable to known risk factors. *Birth Defects Res A Clin Mol Teratol.* 2013;97(1):42-46. doi:10.1002/bdra.23100
  28. Mosley BS, Cleves MA, Siega-Riz AM, et al. Neural tube defects and maternal folate intake among pregnancies conceived after folic acid fortification in the United States. *Am J Epidemiol.* 2009;169(1):9-17. doi:10.1093/aje/kwn331
  29. Czeizel AE, Météneki J, Dudás I. The higher rate of multiple births after periconceptional multivitamin supplementation: an analysis of causes. *Acta Genet Med Gemellol (Roma).* 1994;43(3-4):175-184. doi:10.1017/S0001566000001938
  30. Vollset SE, Gjessing HK, Tandberg A, et al. Folate supplementation and twin pregnancies. *Epidemiology.* 2005;16(2):201-205. doi:10.1097/01.ede.0000152914.84962.13
  31. Bekkers MB, Elstgeest LE, Scholten S, et al. Maternal use of folic acid supplements during pregnancy, and childhood respiratory health and atopy. *Eur Respir J.* 2012;39(6):1468-1474. doi:10.1183/09031936.00094511
  32. Dunstan JA, West C, McCarthy S, et al. The relationship between maternal folate status in pregnancy, cord blood folate levels, and allergic outcomes in early childhood. *Allergy.* 2012;67(1):50-57. doi:10.1111/j.1368-9995.2011.02714.x
  33. Granell R, Heron J, Lewis S, Davey Smith G, Sterne JA, Henderson J. The association between mother and child MTHFR C677T polymorphisms, dietary folate intake and childhood atopy in a population-based, longitudinal birth cohort. *Clin Exp Allergy.* 2008;38(2):320-328. doi:10.1111/j.1365-2222.2007.02902.x
  34. Häberg SE, London SJ, Stigum H, Nafstad P, Nystad W. Folic acid supplements in pregnancy and early childhood respiratory health. *Arch Dis Child.* 2009;94(3):180-184. doi:10.1136/adc.2008.142448
  35. Kieft-de Jong JC, Timmermans S, Jaddoe VW, et al. High circulating folate and vitamin B-12 concentrations in women during pregnancy are associated with increased prevalence of atopic dermatitis in their offspring. *J Nutr.* 2012;142(4):731-738. doi:10.3945/jn.111.154948
  36. Magdelijns FJ, Mommers M, Penders J, Smits L, Thijss C. Folic acid use in pregnancy and the development of atopy, asthma, and lung function in childhood. *Pediatrics.* 2011;128(1):e135-144. doi:10.1542/peds.2010-1690
  37. Martinussen MP, Risnes KR, Jacobsen GW, Bracken MB. Folic acid supplementation in early pregnancy and asthma in children aged 6 years. *Am J Obstet Gynecol.* 2012;206(1):72.e71-77. doi:10.1016/j.ajog.2011.07.033
  38. Whitrow MJ, Moore VM, Rumbold AR, Davies MJ. Effect of supplemental folic acid in pregnancy on childhood asthma: a prospective birth cohort study. *Am J Epidemiol.* 2009;170(12):1486-1493. doi:10.1093/aje/kwp315
  39. Yang L, Jiang L, Bi M, et al. High dose of maternal folic acid supplementation is associated to infant asthma. *Food Chem Toxicol.* 2015;75:88-93. doi:10.1016/j.fct.2014.11.006
  40. Wang T, Zhang HP, Zhang X, Liang ZA, Ji YL, Wang G. Is folate status a risk factor for asthma or other allergic diseases? *Allergy Asthma Immunol Res.* 2015;7(6):538-546. doi:10.4168/aair.2015.7.6.538
  41. Crider KS, Cordero AM, Qi YP, Mulinare J, Dowling NF, Berry RJ. Prenatal folic acid and risk of asthma in children: a systematic review and meta-analysis. *Am J Clin Nutr.* 2013;98(5):1272-1281. doi:10.3945/ajcn.113.065623
  42. Czeizel AE. Periconceptional folic acid containing multivitamin supplementation. *Eur J Obstet Gynecol Reprod Biol.* 1998;78(2):151-161. doi:10.1016/S0301-2115(98)00061-x
  43. Czeizel AE, Dudás I. Prevention of the first occurrence of neural-tube defects by periconceptional vitamin supplementation. *N Engl J Med.* 1992;327(26):1832-1835. doi:10.1056/NEJM199212243272602
  44. Czeizel AE. Prevention of congenital abnormalities by periconceptional multivitamin supplementation. *BMJ.* 1993;306(6893):1645-1648. doi:10.1136/bmj.306.6893.1645
  45. Czeizel AE. Controlled studies of multivitamin supplementation on pregnancy outcomes. *Ann NY Acad Sci.* 1993;678:266-275. doi:10.1111/j.1749-6632.1993.tb26128.x
  46. Czeizel AE. Reduction of urinary tract and cardiovascular defects by periconceptional multivitamin supplementation. *Am J Med Genet.* 1996;62(2):179-183. doi:10.1002/(SICI)1096-8628(19960315)62:2<179::AID-AJMG1>3.0.CO;2-L
  47. Czeizel AE, Dudás I, Météneki J. Pregnancy outcomes in a randomised controlled trial of periconceptional multivitamin supplementation: final report. *Arch Gynecol Obstet.* 1994;255(3):131-139. doi:10.1007/BF02390940